arXiv:1408.5061v2 [math.NT] 9 Apr 2015

CONGRUENCES FOR PARTITION PAIRS WITH CONDITIONS

CHRIS JENNINGS-SHAFFER

Abstract. We prove congruences for the number of partition pairs (π_1, π_2) such that π_1 is non-empty, $s(\pi_1) \leq s(\pi_2)$, and $\ell(\pi_2) < 2s(\pi_1)$ where $s(\pi)$ is the smallest part and $\ell(\pi)$ is the largest part of a partition. The proofs use Bailey's Lemma and a generalized Lambert series identity of Chan. We also discuss how a partition pair crank gives combinatorial refinements of these congruences.

1. Introduction

We recall that a partition of a non-negative integer n is a non-decreasing sequence of positive integers that sum to n. For a partition π we denote the sum of parts by $|\pi|$, the number of parts by $\#(\pi)$, the smallest part by $s(\pi)$, and the largest part by $\ell(\pi)$. We denote the empty partition by $\emptyset$ and use the convention that the empty partition has smallest part ∞ and largest part 0. A partition pair of n is an ordered pair of partitions (π_1, π_2) such that $n = |\pi_1| + |\pi_2|$.

In this paper, we consider the set of partition pairs (π_1, π_2) such that $\pi_1 \neq \emptyset$, $s(\pi_1) \leq s(\pi_2)$, and $\ell(\pi_2) < 2s(\pi_1)$. That is, π_1 is non-empty and each part of π_2 is in the interval $[s(\pi_1), 2s(\pi_1))$. We denote the set of all such partition pairs by $\widetilde{\mathrm{PP}}$. As we will see from the generating function, the number of such partition pairs of n can also be viewed as the number of occurrences of the smallest parts in the partition pairs (π_1, π_2) of n such that $\pi_1 \neq \emptyset$, $s(\pi_1) < s(\pi_2)$, and $\ell(\pi_2) < 2s(\pi_1)$.

We let $\widetilde{\mathrm{pp}}(n)$ denote the number of partition pairs (π_1, π_2) from $\widetilde{\mathrm{PP}}$ of n. As an example, $\widetilde{\mathrm{pp}}(5) = 15$ since the partition pairs are: $(5, \emptyset)$, $(1+4, \emptyset)$, $(2+3, \emptyset)$, $(1+1+3, \emptyset)$, $(1+2+2, \emptyset)$, $(1+1+1+2, \emptyset)$, $(1+1+1+1+1, \emptyset)$, $(1+3, 1)$, $(1+1+2, 1)$, $(1+1+1+1, 1)$, $(1+2, 1+1)$, $(1+1+1, 1+1)$, $(2, 3)$, $(1+1, 1+1+1)$, and $(1, 1+1+1+1)$.

These partition pairs actually came from considering smallest parts partition functions. We recall that $\mathrm{spt}\,(n)$ is the number of occurrences of the smallest part in each partition of n. An overpartion of n is a partition of n in which the first occurrence of a part may be overlined; the number of smallest parts among the overpartitions of n where the smallest part is not overlined is $\overline{\mathrm{spt}}\,(n)$. We use the convention of not including overpartitions when the smallest part is overlined, otherwise if we did include these overpartitions, then we would exactly double $\overline{\mathrm{spt}}\,(n)$. Additionally we have $\overline{\mathrm{spt}}_1\,(n)$ and $\overline{\mathrm{spt}}_2\,(n)$ denoting the restriction of $\overline{\mathrm{spt}}\,(n)$ to overpartitions where the smallest part is odd and even, respectively.

In Section 3 of [8], Garvan and the author considered the partition pairs (π_1, π_2) such that π_1 is non-empty, $s(\pi_1) \leq s(\pi_2)$, and if a part of π_2 is larger than $2s(\pi_1)$ then it must be odd. It turns out the number of such partition pairs of n is $\overline{\mathrm{spt}}\,(n)$. Additionally restricting $s(\pi_1)$ to an odd or even number gives $\overline{\mathrm{spt}}_1\,(n)$ or $\overline{\mathrm{spt}}_2\,(n)$, respectively. The partition pairs here are those were we additionally do not allow the parts of π_2 to be larger than $2s(\pi_1) - 1$.

The smallest parts functions satisfy a wide array of congruences. For example, for $n \geq 0$ we have

$$\mathrm{spt}\,(5n+4) \equiv 0 \pmod 5,$$

2010 *Mathematics Subject Classification.* Primary 11P83, 05A17, 11P82.

Key words and phrases. Andrews' spt-function; congruences; partitions; smallest parts function.

$$\begin{aligned}
\operatorname{spt}(7n+5) &\equiv 0 \pmod 7,\\
\operatorname{spt}(13n+6) &\equiv 0 \pmod{13},\\
\overline{\operatorname{spt}}(3n) &\equiv 0 \pmod 3,\\
\overline{\operatorname{spt}}_1(3n) &\equiv 0 \pmod 3,\\
\overline{\operatorname{spt}}_1(5n) &\equiv 0 \pmod 5,\\
\overline{\operatorname{spt}}_2(3n) &\equiv 0 \pmod 3,\\
\overline{\operatorname{spt}}_2(3n+1) &\equiv 0 \pmod 3,\\
\overline{\operatorname{spt}}_2(5n+3) &\equiv 0 \pmod 5.
\end{aligned}$$

The congruences for $\operatorname{spt}(n)$ were first proved by Andrews in [2] and the congruences for $\overline{\operatorname{spt}}(n)$, $\overline{\operatorname{spt}}_1(n)$, and $\overline{\operatorname{spt}}_2(n)$ were first proved by Bringmann, Lovejoy, and Osburn in [5]. The function $\widetilde{\operatorname{pp}}(n)$ satisfies similar congruences.

Theorem 1.1. *For $n \geq 0$*

$$\begin{aligned}
\widetilde{\operatorname{pp}}(3n+2) &\equiv 0 \pmod 3,\\
\widetilde{\operatorname{pp}}(5n+3) &\equiv 0 \pmod 5,\\
\widetilde{\operatorname{pp}}(5n+4) &\equiv 0 \pmod 5.
\end{aligned}$$

To start, by summing according to the smallest part of π_1 we see a generating function for $\widetilde{\operatorname{pp}}(n)$ is

$$\widetilde{\operatorname{PP}}(q) = \sum_{n=0}^{\infty} \widetilde{\operatorname{pp}}(n) q^n = \sum_{n=1}^{\infty} \frac{q^n}{(q^n;q)_\infty (q^n;q)_n} = \sum_{n=1}^{\infty} \frac{q^n (q^{2n};q)_\infty}{(q^n, q^n;q)_\infty}.$$

By rewriting the summand as $\frac{q^n}{(1-q^n)^2 (q^{n+1};q)_\infty (q^{n+1};q)_{n-1}}$, we see that $\widetilde{\operatorname{pp}}(m)$ is also the number of occurrences of the smallest parts in the partition pairs (π_1, π_2) of m where each part of π_2 is strictly between $s(\pi_1)$ and $2s(\pi_2)$. It is not clear if $\widetilde{\operatorname{pp}}(n)$ can easily be phrased in terms of the smallest parts of any sort of restricted partitions or overpartitions, rather than partition pairs. Here and throughout, we use the following standard q-hypergeometric notation

$$\begin{aligned}
(z;q)_n &= \prod_{k=0}^{n-1} (1 - zq^k),\\
(z;q)_\infty &= \prod_{k=0}^{\infty} (1 - zq^k),\\
(z_1, \ldots, z_n;q)_\infty &= (z_1;q)_\infty \ldots (z_n;q)_\infty,\\
[z;q]_\infty &= (z, q/z;q)_\infty,\\
[z_1, \ldots, z_n;q]_\infty &= [z_1;q]_\infty \ldots [z_n;q]_\infty.
\end{aligned}$$

We consider a two-variable generalization of $\widetilde{\operatorname{PP}}(q)$ given by

$$\widetilde{\operatorname{PP}}(z,q) = \sum_{n=1}^{\infty} \frac{q^n (q^{2n};q)_\infty}{(zq^n, z^{-1}q^n;q)_\infty} = \sum_{n=1}^{\infty} \sum_{m=-\infty}^{\infty} C(m,n) z^m q^n,$$

so that $\widetilde{\operatorname{PP}}(1,q) = \widetilde{\operatorname{PP}}(q)$. We define

$$C(k,t,n) = \sum_{m \equiv k \pmod t} C(m,n),$$

and so for any positive t

$$\widetilde{\mathrm{pp}}(n) = \sum_{m=-\infty}^{\infty} C(m,n) = \sum_{k=0}^{t-1} C(k,t,n). \tag{1.1}$$

We prove Theorem 1.1 with the following strategy. If ζ_t is a primitive t^{th} root of unity with $t = 3$ or 5, then

$$\widetilde{\mathrm{PP}}(\zeta_t, q) = \sum_{n=1}^{\infty} \sum_{m=-\infty}^{\infty} C(m,n)\zeta_t^m q^n = \sum_{n=1}^{\infty} q^n \sum_{k=0}^{t-1} \zeta_t^k C(k,t,n).$$

As the minimal polynomial for ζ_t is $\sum_{k=0}^{t-1} x^k$, if the coefficient of q^N in $\widetilde{\mathrm{PP}}(\zeta_t)$ is zero, then $C(0,t,N) = C(1,t,N) \cdots = C(t-1,t,N)$. Thus, by (1.1), $\widetilde{\mathrm{pp}}(N) = t \cdot C(0,t,N)$ and so $\widetilde{\mathrm{pp}}(N) \equiv 0 \pmod{t}$.

Theorem 1.1 now follows from showing that the coefficient of q^{3n+2} in $\widetilde{\mathrm{PP}}(\zeta_3, q)$ and the coefficient of q^{5n+3} and q^{5n+4} in $\widetilde{\mathrm{PP}}(\zeta_5, q)$ are zero. In stating our theorems, we need to first define the series:

$$\Sigma(z,w,q) = \sum_{n=-\infty}^{\infty} \frac{q^{2n(n+1)} w^n}{1 - zq^n}.$$

For integers a, b, c, we will write

$$\Sigma(a,b,c) = \Sigma(q^a, q^b, q^c).$$

We will prove the following theorems about $\widetilde{\mathrm{PP}}(\zeta_3, q)$ and $\widetilde{\mathrm{PP}}(\zeta_5, q)$.

Theorem 1.2.

$$\widetilde{\mathrm{PP}}(\zeta_3, q) = A_0(q^3) + qA_1(q^3) + q^2 A_2(q^3),$$

where

$$\begin{aligned}
A_0(q) &= \frac{\left(q^9;q^9\right)_\infty \left[q^3;q^9\right]_\infty}{\left[q;q^9\right]_\infty^2 \left[q^4;q^9\right]_\infty} - \frac{1}{\left(q^9;q^9\right)_\infty \left[q^2;q^9\right]_\infty} \left(\Sigma(1,-14,9) + q\Sigma(1,-5,9)\right) \\
&\quad - \frac{1}{\left(q^9;q^9\right)_\infty \left[q^4;q^9\right]_\infty} \left(q\Sigma(1,-8,9) + q^5\Sigma(4,1,9)\right), \\
A_1(q) &= \frac{\left(q^3;q^3\right)_\infty}{\left[q;q^3\right]_\infty}, \\
A_2(q) &= 0.
\end{aligned}$$

Theorem 1.3.

$$\widetilde{\mathrm{PP}}(\zeta_5, q) = B_0(q^5) + qB_1(q^5) + q^2 B_2(q^5) + q^3 B_3(q^5) + q^4 B_4(q^5),$$

where

$$\begin{aligned}
B_0(q) &= -\frac{\zeta_5 + \zeta_5^4}{\left(q^5;q^5\right)_\infty} \left(q^{15}\Sigma(7,10,15) + q^{25}\Sigma(13,25,15)\right), \\
B_1(q) &= -\frac{1}{\left(q^5;q^5\right)_\infty} \left(q^{13}\Sigma(10,10,15) + q^{23}\Sigma(10,25,15)\right) + \frac{\left(q^3;q^3\right)_\infty^3}{\left(q;q\right)_\infty \left(q^5;q^5\right)_\infty}, \\
B_2(q) &= \frac{1 + \zeta_5 + \zeta_5^4}{\left(q^5;q^5\right)_\infty} \left(\Sigma(1,-20,15) + q^4\Sigma(4,-5,15)\right), \\
B_3(q) &= 0,
\end{aligned}$$

$$B_4(q) = 0.$$

The partition pairs from Section 3 of [8], were an intermediate step to defining a crank on the smallest parts of overpartitions. This crank gave a combinatorial interpretation of the congruences for $\overline{\text{spt}}(n)$, $\overline{\text{spt}}_1(n)$, and $\overline{\text{spt}}_2(n)$. In [8] we let $k(\pi_1, \pi_2)$ denote the number of parts of π_2 that are between $s(\pi_1)$ and $2s(\pi_1) - 1$ and defined

$$\overline{\text{crank}}(\pi_1, \pi_2) = \begin{cases} (\#\text{ of parts of } \pi_1) - 1 & \text{if } k(\pi_1, \pi_2) = 0 \\ (\#\text{ of parts of } \pi_1 \geq s(\pi_1) + k(\pi_1, \pi_2)) - k(\pi_1, \pi_2) & \text{if } k(\pi_1, \pi_2) > 0. \end{cases}$$

For the partition pairs in this article, $k(\pi_1, \pi_2)$ is $\#(\pi_2)$ and so we define

$$\text{paircrank}(\pi_1, \pi_2) = \begin{cases} (\#\text{ of parts of } \pi_1) - 1 & \text{if } \#(\pi_2) = 0 \\ (\#\text{ of parts of } \pi_1 \geq s(\pi_1) + \#(\pi_2)) - \#(\pi_2) & \text{if } \#(\pi_2) > 0. \end{cases}$$

This paircrank yields a combinatorial refinement of the congruences for $\widetilde{\text{pp}}(n)$.

Theorem 1.4. (i) *The residue of the* paircrank *mod* 3 *divides the partition pairs from* $\widetilde{\text{PP}}$ *of* $3n + 2$ *into* 3 *equal classes.*

(ii) *The residue of the* paircrank *mod* 5 *divides the partition pairs from* $\widetilde{\text{PP}}$ *of* $5n+3$ *into* 5 *equal classes.*

(iii) *The residue of the* paircrank *mod* 5 *divides the partition pairs from* $\widetilde{\text{PP}}$ *of* $5n+4$ *into* 5 *equal classes.*

In Section 2, we provide an alternative expression for $\widetilde{\text{PP}}(z, q)$. In Sections 3 and 4, we prove Theorems 1.2 and 1.3, respectively. In Section 5, we prove Theorem 1.4 by showing that $C(m, n)$ is the number of partition pairs from $\widetilde{\text{PP}}$ of n with paircrank m.

2. Preliminaries

Before proving Theorems 1.2 and 1.3, we require another form for $\widetilde{\text{PP}}(z, q)$. We recall a pair of sequences (α, β) are a Bailey pair relative to (a, q) if

$$\beta_n = \sum_{k=0}^{n} \frac{\alpha_k}{(q;q)_{n-k} (aq;q)_{n+k}}.$$

A limiting case of Bailey's lemma states that if (α, β) are a Bailey pair for (a, q), then

$$\sum_{n=0}^{\infty} (\rho_1, \rho_2; q)_n \left(\frac{aq}{\rho_1\rho_2}\right)^n \beta_n = \frac{(aq/\rho_1, aq/\rho_2; q)_\infty}{(aq, aq/(\rho_1\rho_2); q)_\infty} \sum_{n=0}^{\infty} \frac{(\rho_1, \rho_2; q)_n \left(\frac{aq}{\rho_1\rho_2}\right)^n \alpha_n}{(aq/\rho_1, aq/\rho_2; q)_n}.$$

Proposition 2.1. *The pair* (α, β) *where*

$$\beta_n = \frac{1}{(q;q)_{2n-1}}, \qquad \alpha_{3n} = 0, \qquad \alpha_{3n\pm1} = q^{6n^2\pm n} - q^{6n^2\pm 7n+2},$$

is a Bailey pair relative to $(1, q)$. *Here* $\alpha_0 = \beta_0 = 0$.

Proof. The following is (2.1) of [13],

$$\sum_{n=-\infty}^{\infty} \frac{(1 - aq^{2n}) (b, c, d, e; q)_n a^{2n} q^n}{(1 - a) (aq/b, aq/c, aq/d, aq/e; q)_n b^n c^n d^n e^n}$$
$$= \frac{(q, q/a, qa, aq/bc, aq/, bd, aq/be, aq/cd, aq/de; q)_\infty}{(q/b, q/c, q/d, q/e, aq/b, aq/c, aq/d, aq/e, a^2q/bcde; q)_\infty}.$$

With $q \mapsto q^3$, $b = q^{1-N}$, $c = q^{2-N}$, $d = q^{3-N}$, $a = q^2$, and $e \to \infty$, this becomes

$$\sum_{n=-\infty}^{\infty} \frac{(1-q^{6n+2})\left(q^{1-N};q\right)_{3n}(-1)^n q^{\frac{3n(n-1)}{2}+n+3nN}}{(1-q^2)\left(q^{2+N};q\right)_{3n}} = \frac{\left(q^3,q,q^5;q^3\right)_\infty \left(q^{2N};q\right)_\infty}{\left(q^N,q^{N+2};q\right)_\infty}.$$

Here the sum is actually finite as we see the summands are eventually zero for large positive and large negative values of n. With this in mind, we have for $N > 0$

$$\begin{aligned}
&\sum_{n=0}^{N} \frac{\alpha_n}{(q;q)_{N+n}(q;q)_{N-n}} \\
&= \frac{\alpha_1}{(q;q)_{N-1}(q;q)_{N+1}} + \sum_{n=1}^{\infty} \frac{\alpha_{3n-1}}{(q;q)_{N-3n+1}(q;q)_{N+3n-1}} + \frac{\alpha_{3n+1}}{(q;q)_{N-3n-1}(q;q)_{N+3n+1}} \\
&= \sum_{n=-\infty}^{\infty} \frac{q^{6n^2+n} - q^{6n^2+7n+2}}{(q;q)_{N-3n-1}(q;q)_{N+3n+1}} \\
&= \sum_{n=-\infty}^{\infty} \frac{q^{6n^2+n}(1-q^{6n+2})\left(q^{1-N};q\right)_{3n}(-1)^n q^{-\binom{3n}{2}+3n(N-1)}}{(q;q)_{N-1}(q;q)_{N+1}\left(q^{N+2};q\right)_{3n}} \\
&= \frac{1}{(q;q)_{N-1}(q;q)_{N+1}} \sum_{n=-\infty}^{\infty} \frac{q^{6n^2+n}(1-q^{6n+2})\left(q^{1-N};q\right)_{3n}(-1)^n q^{\frac{3n(n-1)}{2}+n+3nN}}{\left(q^{N+2};q\right)_{3n}} \\
&= \frac{(1-q^2)}{(q;q)_{N-1}(q;q)_{N+1}} \cdot \frac{\left(q^3,q,q^5;q^3\right)_\infty \left(q^{2N};q\right)_\infty}{\left(q^N,q^{N+2};q\right)_\infty} \\
&= \frac{\left(q,q^{2N};q\right)_\infty}{(q,q;q)_\infty} \\
&= \frac{1}{(q;q)_{2N-1}}.
\end{aligned}$$

Thus, the result follows. □

Corollary 2.2.

$$\widetilde{\mathrm{PP}}(z,q) = \frac{1}{(q;q)_\infty} \sum_{n=-\infty}^{\infty} \frac{q^{6n^2+4n+1}(1-q^{6n+2})}{(1-zq^{3n+1})(1-z^{-1}q^{3n+1})}.$$

Proof. By Proposition 2.1, we have

$$\begin{aligned}
\widetilde{\mathrm{PP}}(z,q) &= \frac{(q;q)_\infty}{(z,z^{-1};q)_\infty} \sum_{n=1}^{\infty} \frac{\left(z,z^{-1};q\right)_n q^n}{(q;q)_{2n-1}} \\
&= \frac{(q;q)_\infty}{(z,z^{-1};q)_\infty} \sum_{n=0}^{\infty} \left(z,z^{-1};q\right)_n q^n \beta_n \\
&= \frac{1}{(1-z)(1-z^{-1})(q;q)_\infty} \sum_{n=0}^{\infty} \frac{\left(z,z^{-1};q\right)_n q^n \alpha_n}{\left(zq,z^{-1}q;q\right)_n} \\
&= \frac{1}{(q;q)_\infty} \sum_{n=0}^{\infty} \frac{q^n \alpha_n}{(1-zq^n)(1-z^{-1}q^n)}
\end{aligned}$$

$$= \frac{1}{(q;q)_\infty} \sum_{n=0}^{\infty} \frac{q^{3n+1}\alpha_{3n+1}}{(1-zq^{3n+1})(1-z^{-1}q^{3n+1})} + \sum_{n=1}^{\infty} \frac{q^{3n-1}\alpha_{3n-1}}{(1-zq^{3n-1})(1-z^{-1}q^{3n-1})}$$
$$= \frac{1}{(q;q)_\infty} \sum_{n=0}^{\infty} \frac{q^{6n^2+4n+1}(1-q^{6n+2})}{(1-zq^{3n+1})(1-z^{-1}q^{3n+1})} + \sum_{n=-\infty}^{-1} \frac{q^{6n^2-2n-1}(1-q^{6n+2})}{(1-zq^{-3n-1})(1-z^{-1}q^{-3n-1})}$$
$$= \frac{1}{(q;q)_\infty} \sum_{n=-\infty}^{\infty} \frac{q^{6n^2+4n+1}(1-q^{6n+2})}{(1-zq^{3n+1})(1-z^{-1}q^{3n+1})}.$$

□

Next we define

$$U_\ell(b) = \sum_{n=-\infty}^{\infty} \frac{q^{6n^2+bn}}{1-q^{\ell(3n+1)}}.$$

We use the additional product notation

$$j(z;q) = [z;q]_\infty (q;q)_\infty = \sum_{n=-\infty}^{\infty} (-1)^n z^n q^{n(n-1)/2}.$$

We now provide properties of $[z;q]_\infty$, $j(z,q)$, and $\Sigma(z,w,q)$. The proofs are routine and thus omitted.

Proposition 2.3.

$$[z;q]_\infty = [q/z;q]_\infty, \tag{2.1}$$
$$[z;q]_\infty = -z[qz;q]_\infty, \tag{2.2}$$
$$[z;q]_\infty = -z\left[z^{-1};q\right]_\infty, \tag{2.3}$$
$$\Sigma(z,w,q) = -z^{-1}\Sigma\left(z^{-1},w^{-1}q^{-3},q\right), \tag{2.4}$$
$$\Sigma(z,w,q) = -z^{-1}w^{-1}q\Sigma\left(z^{-1}q,w^{-1}q,q\right), \tag{2.5}$$
$$\Sigma\left(z,z^4/q^2,q\right) + z\Sigma\left(z,z^4/q,q\right) = z^{-1}\Sigma\left(z,z^4/q^3,q\right) + z^2\Sigma\left(z,z^4,q\right) - z^{-1}j(q/z^2;q). \tag{2.6}$$

3. Proof of Theorem 1.2

Lemma 3.1.

$$\frac{(q;q)_\infty^2}{[b_1,1/b_1,b_3,b_4;q]_\infty}$$
$$= \frac{1}{[1/b_1^2,b_3/b_1,b_4/b_1;q]_\infty}\Sigma\left(b_1,\frac{b_1^4}{b_3b_4},q\right) - \frac{b_1}{[b_1^2,b_3b_1,b_4b_1;q]_\infty}\Sigma\left(b_1,\frac{b_1^4b_3b_4}{q^3},q\right)$$
$$+ \frac{1}{[b_1/b_3,1/b_1b_3,b_4/b_3;q]_\infty}\Sigma\left(b_3,\frac{b_3^3}{b_4},q\right) + \frac{1}{[b_1/b_4,1/b_1b_4,b_3/b_4;q]_\infty}\Sigma\left(b_4,\frac{b_4^3}{b_3},q\right).$$

Proof. We will use the $s=4$, $r=0$ case of Theorem 2.1 in [6]:

$$\frac{(q;q)_\infty^2}{[b_1,b_2,b_3,b_4;q]_\infty}$$
$$= \frac{1}{[b_2/b_1,b_3/b_1,b_4/b_1;q]_\infty}\Sigma\left(b_1,\frac{b_1^3}{b_2b_3b_4},q\right) + \frac{1}{[b_1/b_2,b_3/b_2,b_4/b_2;q]_\infty}\Sigma\left(b_2,\frac{b_2^3}{b_1b_3b_4},q\right)$$

$$+\frac{1}{[b_1/b_3,b_2/b_3,b_4/b_3;q]_\infty}\Sigma\left(b_3,\frac{b_3^3}{b_1b_2b_4},q\right)+\frac{1}{[b_1/b_4,b_2/b_4,b_3/b_4;q]_\infty}\Sigma\left(b_4,\frac{b_4^3}{b_1b_2b_3},q\right).$$

Setting $b_2=b_1^{-1}$ we get

$$\begin{aligned}&\frac{(q;q)_\infty^2}{[b_1,1/b_1,b_3,b_4;q]_\infty}\\&=\frac{1}{[1/b_1^2,b_3/b_1,b_4/b_1;q]_\infty}\Sigma\left(b_1,\frac{b_1^4}{b_3b_4},q\right)+\frac{1}{[b_1^2,b_3b_1,b_4b_1;q]_\infty}\Sigma\left(b_1^{-1},\frac{1}{b_1^4b_3b_4},q\right)\\&\quad+\frac{1}{[b_1/b_3,1/b_1b_3,b_4/b_3;q]_\infty}\Sigma\left(b_3,\frac{b_3^3}{b_4},q\right)+\frac{1}{[b_1/b_4,1/b_1b_4,b_3/b_4;q]_\infty}\Sigma\left(b_4,\frac{b_4^3}{b_3},q\right).\end{aligned}$$

By (2.4)

$$\Sigma\left(b_1^{-1},\frac{1}{b_1^4b_3b_4},q\right)=-b_1\Sigma\left(b_1,\frac{b_1^4b_3b_4}{q^3},q\right),$$

and so the result follows. □

Additionally setting $b_4=b_3^{-1}$ gives the following Lemma.

Lemma 3.2.

$$\begin{aligned}&\frac{(q;q)_\infty^2}{[b_1,1/b_1,b_3,1/b_3;q]_\infty}\\&=\frac{1}{[1/b_1^2,b_3/b_1,1/b_1b_3;q]_\infty}\Sigma\left(b_1,b_1^4,q\right)-\frac{b_1}{[b_1^2,b_3b_1,b_1/b_3;q]_\infty}\Sigma\left(b_1,\frac{b_1^4}{q^3},q\right)\\&\quad+\frac{1}{[b_1/b_3,1/b_1b_3,1/b_3^2;q]_\infty}\Sigma\left(b_3,b_3^4,q\right)-\frac{b_3}{[b_1b_3,b_3/b_1,b_3^2;q]_\infty}\Sigma\left(b_3,\frac{b_3^4}{q^3},q\right).\end{aligned}$$

We now require the following Propositions.

Proposition 3.3.

$$\begin{aligned}&\Sigma(1,-11,9)+q^{15}\Sigma(7,16,9)\\&=\frac{\left(q^9;q^9\right)_\infty^2\left[q^3;q^9\right]_\infty}{[q;q^9]_\infty\left[q^4;q^9\right]_\infty}-\frac{\left[q;q^9\right]_\infty}{\left[q^4;q^9\right]_\infty}\left(q\Sigma(1,-8,9)+q^5\Sigma(4,1,9)\right),\qquad(3.1)\\&q^6\Sigma(4,4,9)+q^{13}\Sigma(7,13,9)\\&=q\frac{\left(q^9;q^9\right)_\infty^2\left[q^3;q^9\right]_\infty}{\left[q^4;q^9\right]_\infty^2}-\frac{\left[q^2;q^9\right]_\infty}{\left[q^4;q^9\right]_\infty}\left(q\Sigma(1,-8,9)+q^5\Sigma(4,1,9)\right).\qquad(3.2)\end{aligned}$$

Proof. For (3.1) we move all $\Sigma(z,w,q)$ terms to one side, divide by $\left[q,q^2,q^3;q^9\right]_\infty$, and use that $\Sigma(4,1,9)=-q^4\Sigma(5,8,9)$ by (2.5). Equation (3.1) is then equivalent to

$$\begin{aligned}\frac{\left(q^9;q^9\right)_\infty^2}{[q,q,q^2,q^4;q^9]_\infty}&=\frac{1}{[q,q^2,q^3;q^9]_\infty}\Sigma(1,-11,9)+\frac{q^{15}}{[q,q^2,q^3;q^9]_\infty}\Sigma(7,16,9)\\&\quad+\frac{q}{[q^2,q^3,q^4;q^9]_\infty}\Sigma(1,-8,9)-\frac{q^9}{[q^2,q^3,q^4;q^9]_\infty}\Sigma(5,8,9).\qquad(3.3)\end{aligned}$$

In Lemma 3.1 we use $q \mapsto q^9$, $b_1 = q, b_3 = q^7, b_4 = q^5$ to get

$$\begin{aligned}\frac{\left(q^9;q^9\right)_\infty^2}{\left[q,q^{-1},q^7,q^5;q^9\right]_\infty} =& \frac{1}{\left[q^{-2},q^6,q^4;q^9\right]_\infty}\Sigma\left(1,-8,9\right) - \frac{q}{\left[q^2,q^8,q^6;q^9\right]_\infty}\Sigma\left(1,-11,9\right)\\ &+ \frac{1}{\left[q^{-6},q^{-8},q^{-2};q^9\right]_\infty}\Sigma\left(7,16,9\right) + \frac{1}{\left[q^{-4},q^{-6},q^2;q^9\right]_\infty}\Sigma\left(5,8,9\right).\end{aligned}$$

Simplifying the products with Proposition 2.3 yields (3.3).

For (3.2) we move all $\Sigma\left(z,w,q\right)$ terms to one side, divide by $q\left[q,q^2,q^3;q^9\right]_\infty$, and use that $\Sigma\left(1,-8,9\right) = -q^{16}\Sigma\left(8,17,9\right)$ by (2.5). Equation (3.2) is then equivalent to

$$\begin{aligned}\frac{\left(q^9;q^9\right)_\infty^2}{\left[q,q^2,q^4,q^4;q^9\right]_\infty} =& -\frac{q^{16}}{\left[q,q^3,q^4;q^9\right]_\infty}\Sigma(8,17,9) + \frac{q^4}{\left[q,q^3,q^4;q^9\right]_\infty}\Sigma(4,1,9)\\ &+ \frac{q^5}{\left[q,q^2,q^3;q^9\right]_\infty}\Sigma(4,4,9) + \frac{q^{12}}{\left[q,q^2,q^3;q^9\right]_\infty}\Sigma(7,13,9).\end{aligned} \tag{3.4}$$

In Lemma 3.1 we use $q \mapsto q^9$, $b_1 = q^4, b_3 = q^8, b_4 = q^7$ to get

$$\begin{aligned}\frac{\left(q^9;q^9\right)_\infty^2}{\left[q^4,q^{-4},q^7,q^8;q^9\right]_\infty} =& \frac{1}{\left[q^{-8},q^4,q^3;q^9\right]_\infty}\Sigma\left(4,1,9\right) - \frac{q^4}{\left[q^8,q^{12},q^{11};q^9\right]_\infty}\Sigma\left(4,4,9\right)\\ &+ \frac{1}{\left[q^{-4},q^{-12},q^{-1};q^9\right]_\infty}\Sigma\left(8,17,9\right) + \frac{1}{\left[q^{-3},q^{-11},q;q^9\right]_\infty}\Sigma\left(7,13,9\right).\end{aligned}$$

Simplifying the products with Proposition 2.3 yields (3.4). □

Proposition 3.4.

$$\begin{aligned}&q^{11}\Sigma(7,10,9) + q^{18}\Sigma(7,19,9)\\ &= \frac{\left(q^9;q^9\right)_\infty^2\left[q^3;q^9\right]_\infty\left[q^4;q^9\right]_\infty}{\left[q;q^9\right]_\infty\left[q^2;q^9\right]_\infty^2} - \frac{\left[q^4;q^9\right]_\infty}{\left[q^2;q^9\right]_\infty}\left(\Sigma(1,-14,9) + q\Sigma(1,-5,9)\right),\end{aligned} \tag{3.5}$$

$$\begin{aligned}&q^3\Sigma(4,-2,9) + q^7\Sigma(4,7,9)\\ &= -\frac{\left(q^9;q^9\right)_\infty^2\left[q^3;q^9\right]_\infty}{\left[q;q^9\right]_\infty\left[q^4;q^9\right]_\infty} + \frac{\left[q;q^9\right]_\infty}{\left[q^2;q^9\right]_\infty}\left(\Sigma(1,-14,9) + q\Sigma(1,-5,9)\right).\end{aligned} \tag{3.6}$$

Proof. For (3.5), we move all $\Sigma\left(z,w,q\right)$ terms to one side and multiply by $\left[q^2;q^9\right]_\infty$, so that (3.5) is equivalent to

$$\begin{aligned}&\frac{\left(q^9;q^9\right)_\infty^2\left[q^3,q^4;q^9\right]_\infty}{\left[q,q^2;q^9\right]_\infty}\\ &= \left[q^2;q^9\right]_\infty\left(q^{11}\Sigma\left(7,10,9\right) + q^{18}\Sigma\left(7,19,9\right)\right) + \left[q^4;q^9\right]_\infty\left(\Sigma\left(1,-14,9\right) + q\Sigma\left(1,-5,9\right)\right).\end{aligned} \tag{3.7}$$

However by (2.6) and the definition of $j(z;q)$, we find that the right hand side of (3.7) is

$$\begin{aligned}&\left[q^2;q^9\right]_\infty\left(q^4\Sigma\left(7,1,9\right) + q^{25}\Sigma\left(7,28,9\right) - q^4 j\left(q^{-5};q^9\right)\right)\\ &+ \left[q^4;q^9\right]_\infty\left(q^{-1}\Sigma\left(1,-23,9\right) + q^2\Sigma\left(1,4,9\right) - q^{-1}j\left(q^7;q^9\right)\right)\\ &= \left[q^2;q^9\right]_\infty\left(q^4\Sigma\left(7,1,9\right) + q^{25}\Sigma\left(7,28,9\right)\right) + \left[q^4;q^9\right]_\infty\left(q^{-1}\Sigma\left(1,-23,9\right) + q^2\Sigma\left(1,4,9\right)\right).\end{aligned}$$

With the above and dividing (3.7) by $\left[q,q^2,q^3,q^4;q^9\right]_\infty$, we have that (3.5) is equivalent to

$$\frac{\left(q^9;q^9\right)_\infty^2}{\left[q,q,q^2,q^2;q^9\right]_\infty} = \frac{q^4}{\left[q,q^3,q^4;q^9\right]_\infty}\Sigma\left(7,1,9\right) + \frac{q^{25}}{\left[q,q^3,q^4;q^9\right]_\infty}\Sigma\left(7,28,9\right)$$

$$+\frac{q^{-1}}{\left[q,q^2,q^3;q^9\right]_\infty}\Sigma\left(1,-23,9\right)+\frac{q^2}{\left[q,q^2,q^3;q^9\right]_\infty}\Sigma\left(1,4,9\right). \tag{3.8}$$

Setting $q\mapsto q^9, b_1=q^7, b_3=q$ in Lemma 3.2 gives

$$\begin{aligned}\frac{\left(q^9;q^9\right)_\infty^2}{\left[q,q^{-1},q^7,q^{-7};q^9\right]_\infty}=&\frac{1}{\left[q^{-14},q^{-6},q^{-8};q^9\right]_\infty}\Sigma\left(7,28,9\right)-\frac{q^7}{\left[q^{14},q^8,q^6;q^9\right]_\infty}\Sigma\left(7,1,9\right)\\&+\frac{1}{\left[q^6,q^{-8},q^{-2};q^9\right]_\infty}\Sigma\left(1,4,9\right)-\frac{q}{\left[q^8,q^{-6},q^2;q^9\right]_\infty}\Sigma\left(1,-23,9\right).\end{aligned}$$

Simplifying the products gives (3.8).

For (3.6), we move all $\Sigma\left(z,w,q\right)$ terms to one side and multiply by $\left[q^2;q^9\right]_\infty$, so that (3.6) is equivalent to

$$\begin{aligned}&\frac{\left(q^9;q^9\right)_\infty^2\left[q^2,q^3;q^9\right]_\infty}{\left[q,q^4;q^9\right]_\infty}\\&=\left[q;q^9\right]_\infty\left(\Sigma\left(1,-14,9\right)+q\Sigma\left(1,-5,9\right)\right)+\left[q^2;q^9\right]_\infty\left(-q^3\Sigma\left(4,-2,9\right)-q^7\Sigma\left(4,7,9\right)\right).\end{aligned}\tag{3.9}$$

However, by (2.6), the right hand side of (3.9) is

$$\begin{aligned}&\left[q;q^9\right]_\infty\left(q^{-1}\Sigma\left(1,-23,9\right)+q^2\Sigma\left(1,4,9\right)-q^{-1}j\left(q^7;q^9\right)\right)\\&+\left[q^2;q^9\right]_\infty\left(-q^{-1}\Sigma\left(4,-11,9\right)-q^{11}\Sigma\left(4,16,9\right)+q^{-1}j\left(q;q^9\right)\right)\\&=\left[q;q^9\right]_\infty\left(q^{-1}\Sigma\left(1,-23,9\right)+q^2\Sigma\left(1,4,9\right)\right)+\left[q^2;q^9\right]_\infty\left(-q^{-1}\Sigma\left(4,-11,9\right)-q^{11}\Sigma\left(4,16,9\right)\right).\end{aligned}$$

With the above and dividing (3.9) by $\left[q,q^2,q^3,q^4;q^9\right]_\infty$, we have that (3.6) is equivalent to

$$\begin{aligned}\frac{\left(q^9;q^9\right)_\infty^2}{\left[q,q,q^4,q^4;q^9\right]_\infty}=&\frac{q^{-1}}{\left[q^2,q^3,q^4;q^9\right]_\infty}\Sigma\left(1,-23,9\right)+\frac{q^2}{\left[q^2,q^3,q^4;q^9\right]_\infty}\Sigma\left(1,4,9\right)\\&-\frac{q^{-1}}{\left[q,q^3,q^4;q^9\right]_\infty}\Sigma\left(4,-11,9\right)-\frac{q^{11}}{\left[q,q^3,q^4;q^9\right]_\infty}\Sigma\left(4,16,9\right).\end{aligned}\tag{3.10}$$

Setting $q\mapsto q^9, b_1=q, b_3=q^4$ in Lemma 3.2 gives

$$\begin{aligned}\frac{\left(q^9;q^9\right)_\infty^2}{\left[q,q^{-1},q^4,q^{-4};q^9\right]_\infty}=&\frac{1}{\left[q^{-2},q^3,q^{-5};q^9\right]_\infty}\Sigma\left(1,4,9\right)-\frac{q}{\left[q^2,q^5,q^{-3};q^9\right]_\infty}\Sigma\left(1,-23,9\right)\\&+\frac{1}{\left[q^{-3},q^{-5},q^{-8};q^9\right]_\infty}\Sigma\left(4,16,9\right)-\frac{q^4}{\left[q^5,q^3,q^8;q^9\right]_\infty}\Sigma\left(4,-11,9\right).\end{aligned}$$

Simplifying the products gives (3.10). □

Next we make use of

$$\left(q;q\right)_\infty=\left(q^{27};q^{27}\right)_\infty\left(\left[q^{12};q^{27}\right]_\infty-q\left[q^6;q^{27}\right]_\infty-q^2\left[q^3;q^{27}\right]_\infty\right), \tag{3.11}$$

which follows from Euler's pentagonal number theorem and the Jacobi triple product identity.

Proposition 3.5.

$$\begin{aligned}&\frac{\left(q^{27};q^{27}\right)_\infty^2\left[q^9;q^{27}\right]_\infty\left[q^{12};q^{27}\right]_\infty}{\left[q^3;q^{27}\right]_\infty\left[q^6;q^{27}\right]_\infty^2}-2q^2\frac{\left(q^{27};q^{27}\right)_\infty^2\left[q^9;q^{27}\right]_\infty}{\left[q^3;q^{27}\right]_\infty\left[q^{12};q^{27}\right]_\infty}-q^4\frac{\left(q^{27};q^{27}\right)_\infty^2\left[q^9;q^{27}\right]_\infty}{\left[q^{12};q^{27}\right]_\infty^2}\\&=\frac{\left(q;q\right)_\infty\left(q^{27};q^{27}\right)_\infty\left[q^9;q^{27}\right]_\infty}{\left[q^3;q^{27}\right]_\infty^2\left[q^{12};q^{27}\right]_\infty}+q\frac{\left(q;q\right)_\infty\left(q^9;q^9\right)_\infty}{\left[q^3;q^9\right]_\infty}.\end{aligned}$$

Proof. Multiplying both sides by $\frac{\left[q^3,q^{12};q^{27}\right]_\infty}{(q^{27};q^{27})_\infty[q^9;q^{27}]_\infty}$, we find this proposition is equivalent to

$$\left(q^{27};q^{27}\right)_\infty\left(\frac{\left[q^{12};q^{27}\right]_\infty^2}{\left[q^6;q^{27}\right]_\infty^2}-2q^2-q^4\frac{\left[q^3;q^{27}\right]_\infty}{\left[q^{12};q^{27}\right]_\infty}\right)=(q;q)_\infty\left(\frac{1}{\left[q^3;q^{27}\right]_\infty}+\frac{q}{\left[q^3;q^{27}\right]_\infty\left[q^6;q^{27}\right]_\infty}\right). \tag{3.12}$$

By (3.11) we see (3.12) reduces to proving

$$\frac{\left[q^{12};q^{27}\right]_\infty^2}{\left[q^6;q^{27}\right]_\infty^2}=\frac{\left[q^{12};q^{27}\right]_\infty}{\left[q^3;q^{27}\right]_\infty}-q^3\frac{\left[q^3;q^{27}\right]_\infty}{\left[q^6;q^{27}\right]_\infty}, \tag{3.13}$$

$$-q^4\frac{\left[q^3;q^{27}\right]_\infty}{\left[q^{12};q^{27}\right]_\infty}=-q\frac{\left[q^6;q^{27}\right]_\infty}{\left[q^3;q^{27}\right]_\infty}+q\frac{\left[q^{12};q^{27}\right]_\infty}{\left[q^6;q^{27}\right]_\infty}. \tag{3.14}$$

However (3.13) follows from multiplying (3.14) by $q\frac{\left[q^6;q^{27}\right]_\infty}{\left[q^{12};q^{27}\right]_\infty}$ and elementary rearrangements. In (3.14) we replace q by $q^{1/3}$ and clear denominators to see we need only prove that

$$q\left[q,q,q^2;q^9\right]_\infty=\left[q^2,q^2,q^4;q^9\right]_\infty-\left[q,q^4,q^4;q^9\right]_\infty. \tag{3.15}$$

However, this follows from the $q\mapsto q^9, x=q, t=y=z=q^2$ case of equation (2.1) from [9], which is an identity of Jacobi:

$$\frac{z}{x}\left[y,x,xt/z,zty;q\right]_\infty=\left[z,t,xty,zy/x;q\right]_\infty-\left[xt,zy,ty,z/x;q\right]_\infty.$$

□

Letting ζ_3 be a primitive third root of unity, we find that

$$\begin{aligned}\widetilde{\mathrm{PP}}(\zeta_3,q)&=\frac{1}{(q;q)_\infty}\sum_{n=-\infty}^{\infty}\frac{q^{6n^2+4n+1}(1-q^{6n+2})(1-q^{3n+1})}{1-q^{9n+3}}\\&=\frac{1}{(q;q)_\infty}\left(qU_3(4)-q^2U_3(7)-q^3U_3(10)+q^4U(13)\right).\end{aligned} \tag{3.16}$$

Using this form of $\widetilde{\mathrm{PP}}(\zeta_3,q)$ in terms of the $U_3(b)$, we proceed in a manner similar to how Atkin and Swinnerton-Dyer in [4] determined rank difference formulas for Dyson's rank of a partition. This was also used to determine crank difference formulas by Ekin in [7] and various rank difference formulas related to overpartitions by Lovejoy and Osburn in [10, 11, 12]. Here the major difference is that in $\Sigma(z,w,q)$ we never use $z=1$ and so we do not have to introduce an extra function to avoid the issue at $n=0$. To begin we note that

$$\begin{aligned}U_3(b)&=\sum_{n=-\infty}^{\infty}\frac{q^{6n^2+bn}}{1-q^{9n+3}}\\&=\sum_{k=0}^{2}\sum_{n=-\infty}^{\infty}\frac{q^{6(3n+k)^2+b(3n+k)}}{1-q^{9(3n+k)+3}}\\&=\sum_{k=0}^{2}q^{6k^2+bk}\sum_{n=-\infty}^{\infty}\frac{q^{54n(n+1)}q^{36nk+3nb-54n}}{1-q^{9k+3}q^{27n}}\\&=\sum_{k=0}^{2}q^{6k^2+bk}\Sigma\left(9k+3,36k+3b-54,27\right).\end{aligned}$$

Thus

$$\begin{aligned}
&qU_3(4)-q^2U_3(7)-q^3U_3(10)+q^4U_3(13)\\
&=q\Sigma(3,-42,27)+q^{11}\Sigma(12,-6,27)+q^{33}\Sigma(21,30,27)-q^2\Sigma(3,-33,27)-q^{15}\Sigma(12,3,27)\\
&\quad-q^{40}\Sigma(21,39,27)-q^3\Sigma(3,-24,27)-q^{19}\Sigma(12,12,27)-q^{47}\Sigma(21,48,27)+q^4\Sigma(3,-15,27)\\
&\quad+q^{54}\Sigma(21,57,27)+q^{23}\Sigma(12,21,27)\\
&=q\Sigma(3,-42,27)+q^4\Sigma(3,-15,27)-q^3\Sigma(3,-24,27)-q^{15}\Sigma(12,3,27)\\
&\quad+q^{33}\Sigma(21,30,27)+q^{54}\Sigma(21,57,27)+q^{23}\Sigma(12,21,27)+q^{11}\Sigma(12,-6,27)\\
&\quad-q^2\Sigma(3,-33,27)-q^{47}\Sigma(21,48,27)-q^{19}\Sigma(12,12,27)-q^{40}\Sigma(21,39,27).
\end{aligned}$$

In the last line we have ordered the terms to apply Propositions 3.3 and 3.4 with $q\mapsto q^3$. With these identities we have that

$$\begin{aligned}
&qU_3(4)-q^2U_3(7)-q^3U_3(10)+q^4U_3(13)\\
&=\big(\Sigma(3,-42,27)+q^3\Sigma(3,-15,27)\big)\left(q+q^2\frac{\left[q^3;q^{27}\right]_\infty}{\left[q^6;q^{27}\right]_\infty}-\frac{\left[q^{12};q^{27}\right]_\infty}{\left[q^6;q^{27}\right]_\infty}\right)\\
&\quad-\big(q^3\Sigma(3,-24,27)+q^{15}\Sigma(12,3,27)\big)\left(1-q\frac{\left[q^6;q^{27}\right]_\infty}{\left[q^{12};q^{27}\right]_\infty}-q^2\frac{\left[q^3;q^{27}\right]_\infty}{\left[q^{12};q^{27}\right]_\infty}\right)\\
&\quad+\frac{\left(q^{27};q^{27}\right)_\infty^2\left[q^9,q^{12};q^{27}\right]_\infty}{\left[q^3,q^6,q^6;q^{27}\right]_\infty}-2q^2\frac{\left(q^{27};q^{27}\right)_\infty^2\left[q^9;q^{27}\right]_\infty}{\left[q^3,q^{12};q^{27}\right]_\infty}-q^4\frac{\left(q^{27};q^{27}\right)_\infty^2\left[q^9;q^{27}\right]_\infty}{\left[q^{12},q^{12};q^{27}\right]_\infty}.
\end{aligned}$$

Next by (3.11) and Proposition 3.5 we have that

$$\begin{aligned}
&qU_3(4)-q^2U_3(7)-q^3U_3(10)+q^4U_3(13)\\
&=-\frac{(q;q)_\infty}{\left(q^{27};q^{27}\right)_\infty\left[q^6;q^{27}\right]_\infty}\big(\Sigma(3,-42,27)+q^3\Sigma(3,-15,27)\big)\\
&\quad-\frac{(q;q)_\infty}{\left(q^{27};q^{27}\right)_\infty\left[q^{12};q^{27}\right]_\infty}\big(q^3\Sigma(3,-24,27)+q^{15}\Sigma(12,3,27)\big)\\
&\quad+\frac{(q;q)_\infty\left(q^{27};q^{27}\right)_\infty\left[q^9;q^{27}\right]_\infty}{\left[q^3;q^{27}\right]_\infty^2\left[q^{12};q^{27}\right]_\infty}+q\frac{(q;q)_\infty\left(q^9;q^9\right)_\infty}{\left[q^3;q^9\right]_\infty}.
\end{aligned}\tag{3.17}$$

Theorem 1.2 now follows by equations (3.16) and (3.17).

4. Proof of Theorem 1.3

We require the following two Lemmas. Both are applications of Theorem 2.1 of [6], namely we take $s=6$, $r=2$, $b_4=b_1^{-1}$, $b_5=b_2^{-1}$, $b_6=b_3^{-1}$ to obtain Lemma 4.1 and take $s=10$, $r=6$, $b_6=b_1^{-1}$, $b_7=b_2^{-1}$, $b_8=b_3^{-1}$, $b_9=b_4^{-1}$, and $b_{10}=b_5^{-1}$ to obtain Lemma 4.2.

Lemma 4.1.

$$\begin{aligned}
\frac{[a_1,a_2;q]_\infty(q;q)_\infty^2}{\left[b_1,b_1^{-1},b_2,b_2^{-1},b_3,b_3^{-1};q\right]_\infty}=&\frac{\left[a_1b_1^{-1},a_2b_1^{-1};q\right]_\infty}{\left[b_1^{-1}b_2,b_1^{-1}b_3,b_1^{-1}b_2^{-1},b_1^{-1}b_3^{-1},b_1^{-2};q\right]_\infty}\Sigma\left(b_1,a_1a_2b_1^4,q\right)\\
&-\frac{b_1\left[a_1b_1,a_2b_1;q\right]_\infty}{\left[b_1b_2,b_1b_3,b_1b_2^{-1},b_1b_3^{-1},b_1^2;q\right]_\infty}\Sigma\left(b_1,\frac{b_1^4}{a_1a_2q^3},q\right)
\end{aligned}$$

$$+\frac{\left[a_1b_2^{-1},a_2b_2^{-1};q\right]_\infty}{\left[b_1b_2^{-1},b_2^{-1}b_3,b_1^{-1}b_2^{-1},b_2^{-1}b_3^{-1},b_2^{-2};q\right]_\infty}\Sigma\left(b_2,a_1a_2b_2^4,q\right)$$
$$-\frac{b_2\left[a_1b_2,a_2b_2;q\right]_\infty}{\left[b_1b_2,b_2b_3,b_1^{-1}b_2,b_2b_3^{-1},b_2^2;q\right]_\infty}\Sigma\left(b_2,\frac{b_2^4}{a_1a_2q^3},q\right)$$
$$+\frac{\left[a_1b_3^{-1},a_2b_3^{-1};q\right]_\infty}{\left[b_1b_3^{-1},b_2b_3^{-1},b_1^{-1}b_3^{-1},b_2^{-1}b_3^{-1},b_3^{-2};q\right]_\infty}\Sigma\left(b_3,a_1a_2b_3^4,q\right)$$
$$-\frac{b_3\left[a_1b_3,a_2b_3;q\right]_\infty}{\left[b_1b_3,b_2b_3,b_1^{-1}b_3,b_2^{-1}b_3,b_3^2;q\right]_\infty}\Sigma\left(b_3,\frac{b_3^4}{a_1a_2q^3},q\right).$$

Lemma 4.2.

$$\frac{\left[a_1,a_2,a_3,a_4,a_5,a_6;q\right]_\infty(q;q)_\infty^2}{\left[b_1,b_1^{-1},b_2,b_2^{-1},b_3,b_3^{-1},b_4,b_4^{-1},b_5,b_5^{-1};q\right]_\infty}$$
$$=\frac{\left[a_1b_1^{-1},a_2b_1^{-1},a_3b_1^{-1},a_4b_1^{-1},a_5b_1^{-1},a_6b_1^{-1};q\right]_\infty}{\left[b_1^{-1}b_2,b_1^{-1}b_3,b_1^{-1}b_4,b_1^{-1}b_5,b_1^{-1}b_2^{-1},b_1^{-1}b_3^{-1},b_1^{-1}b_4^{-1},b_1^{-1}b_5^{-1},b_1^{-2};q\right]_\infty}\Sigma\left(b_1,a_1a_2a_3a_4a_5a_6b_1^4,q\right)$$
$$-\frac{b_1\left[a_1b_1,a_2b_1,a_3b_1,a_4b_1,a_5b_1,a_6b_1;q\right]_\infty}{\left[b_1b_2,b_1b_3,b_1b_4,b_1b_5,b_1b_2^{-1},b_1b_3^{-1},b_1b_4^{-1},b_1b_5^{-1},b_1^2;q\right]_\infty}\Sigma\left(b_1,\frac{b_1^4}{a_1a_2a_3a_4a_5a_6q^3},q\right)$$
$$+\frac{\left[a_1b_2^{-1},a_2b_2^{-1},a_3b_2^{-1},a_4b_2^{-1},a_5b_2^{-1},a_6b_2^{-1};q\right]_\infty}{\left[b_1b_2^{-1},b_2^{-1}b_3,b_2^{-1}b_4,b_2^{-1}b_5,b_1^{-1}b_2^{-1},b_2^{-1}b_3^{-1},b_2^{-1}b_4^{-1},b_2^{-1}b_5^{-1},b_2^{-2};q\right]_\infty}\Sigma\left(b_2,a_1a_2a_3a_4a_5a_6b_3^4,q\right)$$
$$-\frac{b_2\left[a_1b_2,a_2b_2,a_3b_2,a_4b_2,a_5b_2,a_6b_2;q\right]_\infty}{\left[b_1b_2,b_2b_3,b_2b_4,b_2b_5,b_1^{-1}b_2,b_2b_3^{-1},b_2b_4^{-1},b_2b_5^{-1},b_2^2;q\right]_\infty}\Sigma\left(b_2,\frac{b_2^4}{a_1a_2a_3a_4a_5a_6q^3},q\right)$$
$$+\frac{\left[a_1b_3^{-1},a_2b_3^{-1},a_3b_3^{-1},a_4b_3^{-1},a_5b_3^{-1},a_6b_3^{-1};q\right]_\infty}{\left[b_1b_3^{-1},b_2b_3^{-1},b_3^{-1}b_4,b_3^{-1}b_5,b_1^{-1}b_3^{-1},b_2^{-1}b_3^{-1},b_3^{-1}b_4^{-1},b_3^{-1}b_5^{-1},b_3^{-2};q\right]_\infty}\Sigma\left(b_3,a_1a_2a_3a_4a_5a_6b_3^4,q\right)$$
$$-\frac{b_3\left[a_1b_3,a_2b_3,a_3b_3,a_4b_3,a_5b_3,a_6b_3;q\right]_\infty}{\left[b_1b_3,b_2b_3,b_3b_4,b_3b_5,b_1^{-1}b_3,b_2^{-1}b_3,b_3b_4^{-1},b_3b_5^{-1},b_3^2;q\right]_\infty}\Sigma\left(b_3,\frac{b_3^4}{a_1a_2a_3a_4a_5a_6q^3},q\right)$$
$$+\frac{\left[a_1b_4^{-1},a_2b_4^{-1},a_3b_4^{-1},a_4b_4^{-1},a_5b_4^{-1},a_6b_4^{-1};q\right]_\infty}{\left[b_1b_4^{-1},b_2b_4^{-1},b_3b_4^{-1},b_4^{-1}b_5,b_1^{-1}b_4^{-1},b_2^{-1}b_4^{-1},b_3^{-1}b_4^{-1},b_4^{-1}b_5^{-1},b_4^{-2};q\right]_\infty}\Sigma\left(b_4,a_1a_2a_3a_4a_5a_6b_4^4,q\right)$$
$$-\frac{b_4\left[a_1b_4,a_2b_4,a_3b_4,a_4b_4,a_5b_4,a_6b_4;q\right]_\infty}{\left[b_1b_4,b_2b_4,b_3b_4,b_4b_5,b_1^{-1}b_4,b_2^{-1}b_4,b_3^{-1}b_4,b_4b_5^{-1},b_4^2;q\right]_\infty}\Sigma\left(b_4,\frac{b_4^4}{a_1a_2a_3a_4a_5a_6q^3},q\right)$$
$$+\frac{\left[a_1b_5^{-1},a_2b_5^{-1},a_3b_5^{-1},a_4b_5^{-1},a_5b_5^{-1},a_6b_5^{-1};q\right]_\infty}{\left[b_1b_5^{-1},b_2b_5^{-1},b_3b_5^{-1},b_4b_5^{-1},b_1^{-1}b_5^{-1},b_2^{-1}b_5^{-1},b_3^{-1}b_5^{-1},b_4^{-1}b_5^{-1},b_5^{-2};q\right]_\infty}\Sigma\left(b_5,a_1a_2a_3a_4a_5a_6b_5^4,q\right)$$
$$-\frac{b_5\left[a_1b_5,a_2b_5,a_3b_5,a_4b_5,a_5b_5,a_6b_5;q\right]_\infty}{\left[b_1b_5,b_2b_5,b_3b_5,b_4b_5,b_1^{-1}b_5,b_2^{-1}b_5,b_3^{-1}b_5,b_4^{-1}b_5,b_5^2;q\right]_\infty}\Sigma\left(b_5,\frac{b_5^4}{a_1a_2a_3a_4a_5a_6q^3},q\right).$$

In the following Propositions we will repeatedly use that $\left[q,q^4,q^6;q^{15}\right]_\infty=\left[q;q^5\right]_\infty$ and $\left[q^2,q^3,q^7;q^{15}\right]_\infty=\left[q^2;q^5\right]_\infty$ in reducing the products. The proofs of these Propositions are similar to the proofs of Propositions 3.3 and 3.4. By moving the $\Sigma(z,w,q)$ terms all to one side and multiplying by the appropriate product, we find the identities are equivalent to a specialization of Lemma 4.1 or 4.2.

Proposition 4.3.

$$\Sigma(7,-2,15)+q^7\Sigma(7,13,15)+q^{16}\Sigma(13,22,15)+q^{29}\Sigma(13,37,15)$$

$$= \frac{\left[q;q^5\right]_\infty}{\left[q^2;q^5\right]_\infty}\left(q^7\Sigma\left(10,10,15\right)+q^{17}\Sigma\left(10,25,15\right)\right)-q^{-6}\frac{\left(q^3;q^3\right)_\infty^3}{\left(q^5;q^5\right)_\infty\left[q^2;q^5\right]_\infty^2}, \tag{4.1}$$

$$\Sigma\left(1,-26,15\right)+q\Sigma\left(1,-11,15\right)+q^2\Sigma\left(4,-14,15\right)+q^6\Sigma\left(4,1,15\right)$$

$$= -\frac{\left[q^2;q^5\right]_\infty}{\left[q;q^5\right]_\infty}\left(q^{13}\Sigma\left(10,10,15\right)+q^{23}\Sigma\left(10,25,15\right)\right)+\frac{\left(q^3;q^3\right)_\infty^3}{\left(q^5;q^5\right)_\infty\left[q;q^{15}\right]_\infty^2}. \tag{4.2}$$

Proof. In Lemma 4.1 we use $q\mapsto q^{15}$, $a_1=q^{-9}$, $a_2=q^{-21}$, $b_1=q^7$, $b_2=q^{10}$, $b_3=q^{13}$ to get

$$\begin{aligned}
&\frac{\left[q^{-9},q^{-21};q^{15}\right]_\infty\left(q^{15};q^{15}\right)_\infty^2}{\left[q^7,q^{10},q^{13},q^{-7},q^{-10},q^{-13};q^{15}\right]_\infty}\\
&=\frac{\left[q^{-16},q^{-28};q^{15}\right]_\infty}{\left[q^3,q^6,q^{-14},q^{-17},q^{-20};q^{15}\right]_\infty}\Sigma\left(7,-2,15\right)-q^7\frac{\left[q^{-2},q^{-14};q^{15}\right]_\infty}{\left[q^{14},q^{17},q^{20},q^{-3},q^{-6};q^{15}\right]_\infty}\Sigma\left(7,13,15\right)\\
&\quad+\frac{\left[q^{-19},q^{-31};q^{15}\right]_\infty}{\left[q^{-3},q^3,q^{-17},q^{-20},q^{-23};q^{15}\right]_\infty}\Sigma\left(10,10,15\right)-q^{10}\frac{\left[q^{-11};q^{15}\right]_\infty}{\left[q^{17},q^{20},q^{23},q^3,q^{-3};q^{15}\right]_\infty}\Sigma\left(10,25,15\right)\\
&\quad+\frac{\left[q^{-22},q^{-34};q^{15}\right]_\infty}{\left[q^{-6},q^{-3},q^{-20},q^{-23},q^{-26};q^{15}\right]_\infty}\Sigma\left(13,22,15\right)-q^{13}\frac{\left[q^4,q^{-8};q^{15}\right]_\infty}{\left[q^{20},q^{23},q^{26},q^6,q^3;q^{15}\right]_\infty}\Sigma\left(13,37,15\right).
\end{aligned}$$

Simplifying the products yields

$$\begin{aligned}
&q^{-6}\frac{\left[q^6,q^6;q^{15}\right]_\infty\left(q^{15};q^{15}\right)_\infty^2}{\left[q^2,q^2,q^5,q^5,q^7,q^7;q^{15}\right]_\infty}\\
&=-\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(7,-2,15\right)-q^7\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(7,13,15\right)\\
&\quad+q^7\frac{\left[q,q^4;q^{15}\right]_\infty}{\left[q^2,q^3,q^3,q^5,q^7;q^{15}\right]_\infty}\Sigma\left(10,10,15\right)+q^{17}\frac{\left[q,q^4;q^{15}\right]_\infty}{\left[q^2,q^3,q^3,q^7;q^{15}\right]_\infty}\Sigma\left(10,25,15\right)\\
&\quad-q^{16}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(13,22,15\right)-q^{29}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(13,37,15\right).
\end{aligned} \tag{4.3}$$

We see that multiplying both sides of (4.3) by $\left[q^3,q^5,q^6;q^{15}\right]_\infty$ implies (4.1) upon noting that $\frac{\left[q^3,q^6,q^6,q^6;q^{15}\right]_\infty\left(q^{15};q^{15}\right)_\infty^2}{\left[q^2,q^2,q^5,q^7,q^7;q^{15}\right]_\infty}=\frac{\left(q^3;q^3\right)_\infty^3}{\left(q^5;q^5\right)_\infty\left[q^2;q^5\right]_\infty^2}$.

In Lemma 4.1 we use $q\mapsto q^{15}$, $a_1=q^{-12}$, $a_2=q^{-18}$, $b_1=q$, $b_2=q^4$, $b_3=q^{10}$ to get

$$\begin{aligned}
&\frac{\left[q^{-12},q^{-18};q^{15}\right]_\infty\left(q^{15};q^{15}\right)_\infty^2}{\left[q,q^4,q^{10},q^{-1},q^{-4},q^{-10};q^{15}\right]_\infty}\\
&=\frac{\left[q^{-13},q^{-19};q^{15}\right]_\infty}{\left[q^3,q^9,q^{-2},q^{-5},q^{-11};q^{15}\right]_\infty}\Sigma\left(1,-26,15\right)-q\frac{\left[q^{-11},q^{-17};q^{15}\right]_\infty}{\left[q^2,q^5,q^{11},q^{-3},q^{-9};q^{15}\right]_\infty}\Sigma\left(1,-11,15\right)\\
&\quad+\frac{\left[q^{-16},q^{-22};q^{15}\right]_\infty}{\left[q^{-3},q^6,q^{-5},q^{-8},q^{-14};q^{15}\right]_\infty}\Sigma\left(4,-14,15\right)-q^4\frac{\left[q^{-8},q^{-14};q^{15}\right]_\infty}{\left[q^5,q^8,q^{14},q^3,q^{-6};q^{15}\right]_\infty}\Sigma\left(4,1,15\right)\\
&\quad+\frac{\left[q^{-22},q^{-28};q^{15}\right]_\infty}{\left[q^{-9},q^{-6},q^{-11},q^{-14},q^{-20};q^{15}\right]_\infty}\Sigma\left(10,10,15\right)-q^{10}\frac{\left[q^{-2},q^{-8};q^{15}\right]_\infty}{\left[q^{11},q^{14},q^{20},q^9,q^6;q^{15}\right]_\infty}\Sigma\left(10,25,15\right).
\end{aligned}$$

Simplifying the products yields

$$
\begin{aligned}
&q^{-18}\frac{\left[q^3,q^3;q^{15}\right]_\infty\left(q^{15};q^{15}\right)_\infty^2}{\left[q,q,q^4,q^4,q^5,q^5;q^{15}\right]_\infty}\\
&=q^{-18}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(1,-26,15\right)+q^{-17}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(1,-11,15\right)\\
&\quad+q^{-16}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(4,-14,15\right)+q^{-12}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(4,1,15\right)\\
&\quad+q^{-5}\frac{\left[q^2,q^7;q^{15}\right]_\infty}{\left[q,q^4,q^5,q^6,q^6;q^{15}\right]_\infty}\Sigma\left(10,10,15\right)+q^5\frac{\left[q^2,q^7;q^{15}\right]_\infty}{\left[q,q^4,q^5,q^6,q^6;q^{15}\right]_\infty}\Sigma\left(10,25,15\right).
\end{aligned}
\tag{4.4}
$$

We see that multiplying both sides of (4.4) by $q^{18}\left[q^3,q^5,q^6;q^{15}\right]_\infty$ implies (4.1) upon noting that $\frac{\left[q^3,q^3,q^3,q^6;q^{15}\right]_\infty\left(q^{15};q^{15}\right)_\infty^2}{\left[q,q,q^4,q^4,q^5;q^{15}\right]_\infty}=\frac{\left(q^3;q^3\right)_\infty^3}{\left(q^5;q^5\right)_\infty\left[q;q^5\right]_\infty^2}$. □

Proposition 4.4.

$$
\begin{aligned}
&\Sigma\left(1,-17,15\right)+q^3\Sigma\left(4,-8,15\right)+q^{10}\Sigma\left(7,7,15\right)+q^{27}\Sigma\left(13,28,15\right)\\
&=\frac{\left[q;q^5\right]_\infty}{\left[q^2;q^5\right]_\infty}\left(\Sigma\left(1,-20,15\right)+q^4\Sigma\left(4,-5,15\right)\right),
\end{aligned}
\tag{4.5}
$$

$$
\begin{aligned}
&\Sigma\left(7,4,15\right)+q^8\Sigma\left(10,16,15\right)+q^{10}\Sigma\left(10,19,15\right)+q^{21}\Sigma\left(13,31,15\right)\\
&=-\frac{\left[q^2;q^5\right]_\infty}{\left[q;q^5\right]_\infty}\left(q^{-9}\Sigma\left(1,-20,15\right)+q^{-5}\Sigma\left(4,-5,15\right)\right)+q^{-9}\frac{\left(q^3;q^3\right)_\infty^3}{\left(q;q\right)_\infty}.
\end{aligned}
\tag{4.6}
$$

Proof. In Lemma 4.2 we use $q\mapsto q^{15}$, $a_1=q^{-15}$, $a_2=q^{-13}$, $a_3=q^{-10}$, $a_4=q^{-8}$, $a_5=q^{10}$, $a_6=q^{12}$, $b_1=q$, $b_2=q^4$, $b_3=q^7$, $b_4=q^{10}$, $b_5=q^{13}$ and note both the product on the left and four terms on the right in Lemma 4.2 are immediately zero, yielding

$$
\begin{aligned}
0=&\frac{\left[q^{-16},q^{-14},q^{-11},q^{-9},q^9,q^{11};q^{15}\right]_\infty}{\left[q^3,q^6,q^9,q^{12},q^{-2},q^{-5},q^{-8},q^{-11},q^{-14};q^{15}\right]_\infty}\Sigma\left(1,-20,15\right)\\
&-q\frac{\left[q^{-14},q^{-12},q^{-9},q^{-7},q^{11},q^{13};q^{15}\right]_\infty}{\left[q^2,q^5,q^8,q^{11},q^{14},q^{-3},q^{-6},q^{-9},q^{-12};q^{15}\right]_\infty}\Sigma\left(1,-17,15\right)\\
&+\frac{\left[q^{-19},q^{-17},q^{-14},q^{-12},q^6,q^8;q^{15}\right]_\infty}{\left[q^{-3},q^3,q^6,q^9,q^{-5},q^{-8},q^{-11},q^{-14},q^{-17};q^{15}\right]_\infty}\Sigma\left(4,-8,15\right)\\
&-q^4\frac{\left[q^{-11},q^{-9},q^{-6},q^{-4},q^{14},q^{16};q^{15}\right]_\infty}{\left[q^5,q^8,q^{11},q^{14},q^{17},q^3,q^{-3},q^{-6},q^{-9};q^{15}\right]_\infty}\Sigma\left(4,-5,15\right)\\
&-q^7\frac{\left[q^{-8},q^{-6},q^{-3},q^{-1},q^{17},q^{19};q^{15}\right]_\infty}{\left[q^8,q^{11},q^{14},q^{17},q^{20},q^6,q^3,q^{-3},q^{-6};q^{15}\right]_\infty}\Sigma\left(7,7,15\right)\\
&+\frac{\left[q^{-28},q^{-26},q^{-23},q^{-21},q^{-3},q^{-1};q^{15}\right]_\infty}{\left[q^{-12},q^{-9},q^{-6},q^{-3},q^{-14},q^{-17},q^{-20},q^{-23},q^{-26};q^{15}\right]_\infty}\Sigma\left(13,28,15\right).
\end{aligned}
$$

Simplifying the products yields

$$
\begin{aligned}
0=&q^{-11}\frac{\left[q,q^4;q^{15}\right]_\infty}{\left[q^2,q^3,q^3,q^5,q^7;q^{15}\right]_\infty}\Sigma\left(1,-20,15\right)-q^{-11}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(1,-17,15\right)\\
&-q^{-8}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(4,-8,15\right)+q^{-7}\frac{\left[q,q^4;q^{15}\right]_\infty}{\left[q^2,q^3,q^3,q^5,q^7;q^{15}\right]_\infty}\Sigma\left(4,-5,15\right)
\end{aligned}
$$

$$- q^{-1}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(7,7,15\right) - q^{16}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(13,28,15\right). \tag{4.7}$$

We see multiplying both sides of (4.7) by $q^{11}\left[q^3,q^5,q^6;q^{15}\right]_\infty$ implies (4.5).

In Lemma 4.2 we use $q \mapsto q^{15}$, $a_1 = q^{-12}$, $a_2 = q^{-11}$, $a_3 = q^{-7}$, $a_4 = q^{-6}$, $a_5 = q^{-2}$, $a_6 = q^{14}$, $b_1 = q$, $b_2 = q^4$, $b_3 = q^7$, $b_4 = q^{10}$, $b_5 = q^{13}$ and note four terms on the right in Lemma 4.2 are immediately zero, yielding

$$\begin{aligned}
&\frac{\left[q^{-12},q^{-11},q^{-7},q^{-6},q^{-2},q^{14};q^{15}\right]_\infty\left(q^{15};q^{15}\right)_\infty}{\left[q^1,q^4,q^7,q^{10},q^{13},q^{-1},q^{-4},q^{-7},q^{-10},q^{-13};q^{15}\right]_\infty}\\
&= \frac{\left[q^{-13},q^{-12},q^{-8},q^{-7},q^{-3},q^{13};q^{15}\right]_\infty}{\left[q^3,q^6,q^9,q^{12},q^{-2},q^{-5},q^{-8},q^{-11},q^{-14};q^{15}\right]_\infty}\Sigma\left(1,-20,15\right)\\
&\quad - q^4\frac{\left[q^{-8},q^{-7},q^{-3},q^{-2},q^2,q^{18};q^{15}\right]_\infty}{\left[q^5,q^8,q^{11},q^{14},q^{17},q^3,q^{-3},q^{-6},q^{-9};q^{15}\right]_\infty}\Sigma\left(4,-5,15\right)\\
&\quad + \frac{\left[q^{-19},q^{-18},q^{-14},q^{-13},q^{-9},q^7;q^{15}\right]_\infty}{\left[q^{-6},q^{-3},q^3,q^6,q^{-8},q^{-11},q^{-14},q^{-17},q^{-20};q^{15}\right]_\infty}\Sigma\left(7,4,15\right)\\
&\quad + \frac{\left[q^{-22},q^{-21},q^{-17},q^{-16},q^{-12},q^4;q^{15}\right]_\infty}{\left[q^{-9},q^{-6},q^{-3},q^3,q^{-11},q^{-14},q^{-17},q^{-20},q^{-23};q^{15}\right]_\infty}\Sigma\left(10,16,15\right)\\
&\quad - q^{10}\frac{\left[q^{-2},q^{-1},q^3,q^4,q^8,q^{24};q^{15}\right]_\infty}{\left[q^{11},q^{14},q^{17},q^{20},q^{23},q^9,q^6,q^3,q^{-3};q^{15}\right]_\infty}\Sigma\left(10,19,15\right)\\
&\quad - q^{13}\frac{\left[q,q^2,q^6,q^7,q^{11},q^{27};q^{15}\right]_\infty}{\left[q^{14},q^{17},q^{20},q^{23},q^{26},q^{12},q^9,q^6,q^3;q^{15}\right]_\infty}\Sigma\left(13,31,15\right).
\end{aligned}$$

Simplifying the products yields

$$\begin{aligned}
&q^{-3}\frac{\left[q^3,q^6;q^{15}\right]_\infty\left(q^{15};q^{15}\right)_\infty^2}{\left[q,q^2,q^4,q^5,q^5,q^7;q^{15}\right]_\infty}\\
&= q^{-3}\frac{\left[q^2,q^7;q^{15}\right]_\infty}{\left[q^1,q^4,q^5,q^6,q^6;q^{15}\right]_\infty}\Sigma\left(1,-20,15\right) + q\frac{\left[q^2,q^7;q^{15}\right]_\infty}{\left[q^1,q^4,q^5,q^6,q^6;q^{15}\right]_\infty}\Sigma\left(4,-5,15\right)\\
&\quad + q^6\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(7,4,15\right) + q^{14}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(10,16,15\right)\\
&\quad + q^{16}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(10,19,15\right) + q^{27}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(13,31,15\right).
\end{aligned} \tag{4.8}$$

We see multiplying (4.8) by $q^{-6}\left[q^3,q^5,q^6;q^{15}\right]_\infty$ implies (4.6), upon noting that $\frac{\left[q^3,q^3,q^6,q^6;q^{15}\right]_\infty\left(q^{15};q^{15}\right)_\infty^2}{\left[q^1,q^2,q^4,q^5,q^7;q^{15}\right]_\infty} = \frac{\left(q^3;q^3\right)_\infty^3}{\left(q;q\right)_\infty}$.

□

Proposition 4.5.

$$\begin{aligned}
&\Sigma\left(1,-14,15\right) + q^2\Sigma\left(4,-11,15\right) + q^7\Sigma\left(7,1,15\right) + q^{32}\Sigma\left(13,34,15\right)\\
&= -\frac{\left[q^2;q^5\right]_\infty}{\left[q;q^5\right]_\infty}\left(q^{11}\Sigma\left(7,10,15\right) + q^{24}\Sigma\left(13,25,15\right)\right),\\
&\Sigma\left(1,-23,15\right) + q^5\Sigma\left(4,-2,15\right) + q^{15}\Sigma\left(10,13,15\right) + q^{21}\Sigma\left(10,22,15\right)
\end{aligned} \tag{4.9}$$

$$= \frac{\left[q;q^5\right]_\infty}{\left[q^2;q^5\right]_\infty}\left(q^{12}\Sigma\left(7,10,15\right)+q^{25}\Sigma\left(13,25,15\right)\right)+\frac{\left(q^3;q^3\right)_\infty^3}{\left(q;q\right)_\infty}. \tag{4.10}$$

Proof. In Lemma 4.2 we use $q\mapsto q^{15}$, $a_1=q^{-15}$, $a_2=q^{-14}$, $a_3=q^{-10}$, $a_4=q^{-9}$, $a_5=q^{10}$, $a_6=q^{11}$, $b_1=q$, $b_2=q^4$, $b_3=q^7$, $b_4=q^{10}$, $b_5=q^{13}$ and note both the product on the left and four terms on the right in Lemma 4.2 are immediately zero, yielding

$$\begin{aligned}
0=&-q\frac{\left[q^{-14},q^{-13},q^{-9},q^{-8},q^{11},q^{12};q^{15}\right]_\infty}{\left[q^2,q^5,q^8,q^{11},q^{14},q^{-3},q^{-6},q^{-9},q^{-12};q^{15}\right]_\infty}\Sigma\left(1,-14,15\right)\\
&+\frac{\left[q^{-19},q^{-18},q^{-14},q^{-13},q^6,q^7;q^{15}\right]_\infty}{\left[q^{-3},q^3,q^6,q^9,q^{-5},q^{-8},q^{-11},q^{-14},q^{-17};q^{15}\right]_\infty}\Sigma\left(4,-11,15\right)\\
&+\frac{\left[q^{-22},q^{-21},q^{-17},q^{-16},q^3,q^4;q^{15}\right]_\infty}{\left[q^{-6},q^{-3},q^3,q^6,q^{-8},q^{-11},q^{-14},q^{-17},q^{-20};q^{15}\right]_\infty}\Sigma\left(7,1,15\right)\\
&-q^7\frac{\left[q^{-8},q^{-7},q^{-3},q^{-2},q^{17},q^{18};q^{15}\right]_\infty}{\left[q^8,q^{11},q^{14},q^{17},q^{20},q^6,q^3,q^{-3},q^{-6};q^{15}\right]_\infty}\Sigma\left(7,10,15\right)\\
&+\frac{\left[q^{-28},q^{-27},q^{-23},q^{-22},q^{-3},q^{-2};q^{15}\right]_\infty}{\left[q^{-12},q^{-9},q^{-6},q^{-3},q^{-14},q^{-17},q^{-20},q^{-23},q^{-26};q^{15}\right]_\infty}\Sigma\left(13,25,15\right)\\
&-q^{13}\frac{\left[q^{-2},q^{-1},q^3,q^4,q^{23},q^{24};q^{15}\right]_\infty}{\left[q^{14},q^{17},q^{20},q^{23},q^{26},q^{12},q^9,q^6,q^3;q^{15}\right]_\infty}\Sigma\left(13,34,15\right).
\end{aligned}$$

Simplifying the products yields

$$\begin{aligned}
0=&-q^{-13}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(1,-14,15\right)-q^{-11}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(4,-11,15\right)\\
&-q^{-6}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(7,1,15\right)-q^{-2}\frac{\left[q^2,q^7;q^{15}\right]_\infty}{\left[q,q^4,q^5,q^6,q^6;q^{15}\right]_\infty}\Sigma\left(7,10,15\right)\\
&-q^{11}\frac{\left[q^2,q^7;q^{15}\right]_\infty}{\left[q,q^4,q^5,q^6,q^6;q^{15}\right]_\infty}\Sigma\left(13,25,15\right)-q^{19}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(13,34,15\right).
\end{aligned} \tag{4.11}$$

We see multiplying (4.11) by $q^{13}\left[q^3,q^5,q^6;q^{15}\right]_\infty$ implies (4.9).

In Lemma 4.2 we use $q\mapsto q^{15}$, $a_1=q^{-13}$, $a_2=q^{-11}$, $a_3=q^{-8}$, $a_4=q^{-6}$, $a_5=q^{-3}$, $a_6=q^{14}$, $b_1=q$, $b_2=q^4$, $b_3=q^7$, $b_4=q^{10}$, $b_5=q^{13}$ and note four terms on the right in Lemma 4.2 are immediately zero, yielding

$$\begin{aligned}
&\frac{\left[q^{-13},q^{-11},q^{-8},q^{-6},q^{-3},q^{14};q^{15}\right]_\infty\left(q^{15};q^{15}\right)_\infty}{\left[q,q^4,q^7,q^{10},q^{13},q^{-1},q^{-4},q^{-7},q^{-10},q^{-13};q^{15}\right]_\infty}\\
&=\frac{\left[q^{-14},q^{-12},q^{-9},q^{-7},q^{-4},q^{13};q^{15}\right]_\infty}{\left[q^3,q^6,q^9,q^{12},q^{-2},q^{-5},q^{-8},q^{-11},q^{-14};q^{15}\right]_\infty}\Sigma\left(1,-23,15\right)\\
&\quad-q^4\frac{\left[q^{-9},q^{-7},q^{-4},q^{-2},q,q^{18};q^{15}\right]_\infty}{\left[q^5,q^8,q^{11},q^{14},q^{17},q^3,q^{-3},q^{-6},q^{-9};q^{15}\right]_\infty}\Sigma\left(4,-2,15\right)\\
&\quad-q^7\frac{\left[q^{-6},q^{-4},q^{-1},q,q^4,q^{21};q^{15}\right]_\infty}{\left[q^8,q^{11},q^{14},q^{17},q^{20},q^6,q^3,q^{-3},q^{-6};q^{15}\right]_\infty}\Sigma\left(7,10,15\right)\\
&\quad+\frac{\left[q^{-23},q^{-21},q^{-18},q^{-16},q^{-13},q^4;q^{15}\right]_\infty}{\left[q^{-9},q^{-6},q^{-3},q^3,q^{-11},q^{-14},q^{-17},q^{-20},q^{-23};q^{15}\right]_\infty}\Sigma\left(10,13,15\right)
\end{aligned}$$

$$- q^{10}\frac{\left[q^{-3},q^{-1},q^2,q^4,q^7,q^{24};q^{15}\right]_\infty}{\left[q^{11},q^{14},q^{17},q^{20},q^{23},q^9,q^6,q^3,q^{-3};q^{15}\right]_\infty}\Sigma\left(10,22,15\right)$$
$$+\frac{\left[q^{-26},q^{-24},q^{-21},q^{-19},q^{-16},q;q^{15}\right]_\infty}{\left[q^{-12},q^{-9},q^{-6},q^{-3},q^{-14},q^{-17},q^{-20},q^{-23},q^{-26};q^{15}\right]_\infty}\Sigma\left(13,25,15\right).$$

Simplifying the products yields

$$\begin{aligned}
&q^{-6}\frac{\left[q^3,q^6;q^{15}\right]_\infty\left(q^{15};q^{15}\right)_\infty^2}{\left[q,q^2,q^4,q^5,q^5,q^7;q^{15}\right]_\infty}\\
&=q^{-6}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(1,-23,15\right)+q^{-1}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(4,-2,15\right)\\
&\quad-q^6\frac{\left[q,q^4;q^{15}\right]_\infty}{\left[q^2,q^3,q^3,q^5,q^7;q^{15}\right]_\infty}\Sigma\left(7,10,15\right)+q^9\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(10,13,15\right)\\
&\quad+q^{15}\frac{1}{\left[q^3,q^5,q^6;q^{15}\right]_\infty}\Sigma\left(10,22,15\right)-q^{19}\frac{\left[q,q^4;q^{15}\right]_\infty}{\left[q^2,q^3,q^3,q^5,q^7;q^{15}\right]_\infty}\Sigma\left(13,25,15\right).
\end{aligned}\tag{4.12}$$

We see multiplying (4.12) by $q^6\left[q^3,q^5,q^6;q^{15}\right]_\infty$ implies (4.10), again noting that $\frac{\left[q^3,q^3,q^6,q^6;q^{15}\right]_\infty\left(q^{15};q^{15}\right)_\infty^2}{\left[q^1,q^2,q^4,q^5,q^7;q^{15}\right]_\infty}=\frac{\left(q^3;q^3\right)_\infty^3}{\left(q;q\right)_\infty}$.

□

Similar to the proof of Theorem 1.2, we begin with expressing $\widetilde{\mathrm{PP}}(\zeta_5,q)$ in terms of $U_5(b)$. With ζ_5 a primitive fifth root of unity, we have

$$\begin{aligned}
\widetilde{\mathrm{PP}}(\zeta_5,q)&=\frac{1}{\left(q;q\right)_\infty}\sum_{n=-\infty}^{\infty}\frac{q^{6n^2+4n+1}(1-q^{6n+2})(1-q^{3n+1})(1-\zeta_5^2q^{3n+1})(1-\zeta_5^3q^{3n+1})}{1-q^{15n+5}}\\
&=\frac{1}{\left(q;q\right)_\infty}\left(qU_5(4)+(\zeta_5+\zeta_5^4)q^2U_5(7)-(1+\zeta_5+\zeta_5^4)q^3U_5(10)-(1+\zeta_5+\zeta_5^4)q^4U_5(13)\right.\\
&\qquad\left.+(\zeta_5+\zeta_5^4)q^5U_5(16)+q^6U_5(19)\right).
\end{aligned}\tag{4.13}$$

We find that

$$\begin{aligned}
U_5(b)&=\sum_{n=-\infty}^{\infty}\frac{q^{6n^2+bn}}{1-q^{15n+5}}\\
&=\sum_{k=0}^{4}q^{6k^2+bk}\sum_{n=-\infty}^{\infty}\frac{q^{150n(n+1)}q^{60nk+5bn-150n}}{1-q^{15k+5}q^{75n}}\\
&=\sum_{k=0}^{4}q^{6k^2+bk}\sum_{n=-\infty}^{\infty}\Sigma\left(15k+5,60k+5b-150,75\right).
\end{aligned}$$

Thus

$$\begin{aligned}
&qU_5(4)-q^3U_5(10)-q^4U_5(13)+q^6U_5(19)\\
&=q\Sigma(5,-130,75)+q^{11}\Sigma(20,-70,75)+q^{33}\Sigma(35,-10,75)+q^{67}\Sigma(50,50,75)+q^{113}\Sigma(65,110,75)\\
&\quad-q^3\Sigma(5,-100,75)-q^{19}\Sigma(20,-40,75)-q^{47}\Sigma(35,20,75)-q^{87}\Sigma(50,80,75)-q^{139}\Sigma(65,140,75)\\
&\quad-q^4\Sigma(5,-85,75)-q^{23}\Sigma(20,-25,75)-q^{54}\Sigma(35,35,75)-q^{97}\Sigma(50,95,75)-q^{152}\Sigma(65,155,75)\\
&\quad+q^6\Sigma(5,-55,75)+q^{31}\Sigma(20,5,75)+q^{68}\Sigma(35,65,75)+q^{117}\Sigma(50,125,75)+q^{178}\Sigma(65,185,75),
\end{aligned}$$

and

$$
\begin{aligned}
&q^2U_5(7) - q^3U_5(10) - q^4U_5(13) + q^5U_5(16)\\
&= q^2\Sigma(5,-115,75) + q^{15}\Sigma(20,-55,75) + q^{40}\Sigma(35,5,75) + q^{77}\Sigma(50,65,75) + q^{126}\Sigma(65,125,75)\\
&\quad - q^3\Sigma(5,-100,75) - q^{19}\Sigma(20,-40,75) - q^{47}\Sigma(35,20,75) - q^{87}\Sigma(50,80,75) - q^{139}\Sigma(65,140,75)\\
&\quad - q^4\Sigma(5,-85,75) - q^{23}\Sigma(20,-25,75) - q^{54}\Sigma(35,35,75) - q^{97}\Sigma(50,95,75) - q^{152}\Sigma(65,155,75)\\
&\quad + q^5\Sigma(5,-70,75) + q^{27}\Sigma(20,-10,75) + q^{61}\Sigma(35,50,75) + q^{107}\Sigma(50,110,75) + q^{165}\Sigma(65,170,75).
\end{aligned}
$$

Next we reorder the $\Sigma(z,w,q)$ terms and apply Propositions 4.3 and 4.4 with $q \mapsto q^5$ to get that

$$
\begin{aligned}
&qU_5(4) - q^3U_5(10) - q^4U_5(13) + q^6U_5(19)\\
&= q^{67}\Sigma(50,50,75) + q^{117}\Sigma(50,125,75) - q^3\Sigma(5,-100,75) - q^{23}\Sigma(20,-25,75)\\
&\quad + q^{33}\left(\Sigma(35,-10,75) + q^{35}\Sigma(35,65,75) + q^{80}\Sigma(65,110,75) + q^{145}\Sigma(65,185,75)\right)\\
&\quad + q\left(\Sigma(5,-130,75) + q^5\Sigma(5,-55,75) + q^{10}\Sigma(20,-70,75) + q^{30}\Sigma(20,5,75)\right)\\
&\quad - q^4\left(\Sigma(5,-85,75) + q^{15}\Sigma(20,-40,75) + q^{50}\Sigma(35,35,75) + q^{135}\Sigma(65,140,75)\right)\\
&\quad - q^{47}\left(\Sigma(35,20,75) + q^{40}\Sigma(50,80,75) + q^{50}\Sigma(50,95,75) + q^{105}\Sigma(65,155,75)\right)\\
&= \left(q^{66}\Sigma(50,50,75) + q^{116}\Sigma(50,125,75)\right)\left(-\frac{\left[q^{10};q^{15}\right]_\infty}{\left[q^5;q^{15}\right]_\infty} + q + q^2\frac{\left[q^5;q^{15}\right]_\infty}{\left[q^{10};q^{15}\right]_\infty}\right)\\
&\quad - \left(q^2\Sigma(5,-100,75) + q^{22}\Sigma(20,-25,75)\right)\left(-\frac{\left[q^{10};q^{15}\right]_\infty}{\left[q^5;q^{15}\right]_\infty} + q + q^2\frac{\left[q^5;q^{15}\right]_\infty}{\left[q^{10};q^{15}\right]_\infty}\right)\\
&\quad - q^3\frac{\left(q^{15};q^{15}\right)_\infty^3}{\left(q^{25};q^{25}\right)_\infty\left[q^{10};q^{25}\right]_\infty^2} + q\frac{\left(q^{15};q^{15}\right)_\infty^3}{\left(q^{25};q^{25}\right)_\infty\left[q^5;q^{25}\right]_\infty^2} - q^2\frac{\left(q^{15};q^{15}\right)_\infty^3}{\left(q^5;q^5\right)_\infty}\\
&= \left(q^{66}\Sigma(50,50,75) + q^{116}\Sigma(50,125,75)\right)\left(-\frac{\left[q^{10};q^{15}\right]_\infty}{\left[q^5;q^{15}\right]_\infty} + q + q^2\frac{\left[q^5;q^{15}\right]_\infty}{\left[q^{10};q^{15}\right]_\infty}\right)\\
&\quad - \left(q^2\Sigma(5,-100,75) + q^{22}\Sigma(20,-25,75)\right)\left(-\frac{\left[q^{10};q^{15}\right]_\infty}{\left[q^5;q^{15}\right]_\infty} + q + q^2\frac{\left[q^5;q^{15}\right]_\infty}{\left[q^{10};q^{15}\right]_\infty}\right)\\
&\quad + q\frac{\left(q^{15};q^{15}\right)_\infty^3}{\left(q^5;q^5\right)_\infty}\left(\frac{\left[q^{10};q^{15}\right]_\infty}{\left[q^5;q^{15}\right]_\infty} - q - q^2\frac{\left[q^5;q^{15}\right]_\infty}{\left[q^{10};q^{15}\right]_\infty}\right).
\end{aligned}
\tag{4.14}
$$

However, by Euler's pentagonal number theorem and the Jacobi triple product identity, we have that

$$
(q;q)_\infty = \left(q^{25};q^{25}\right)_\infty\left(\frac{\left[q^{10};q^{25}\right]_\infty}{\left[q^5;q^{25}\right]_\infty} - q - q^2\frac{\left[q^5;q^{25}\right]_\infty}{\left[q^{10};q^{25}\right]_\infty}\right).
$$

Thus,

$$
\begin{aligned}
&qU_5(4) - q^3U_5(10) - q^4U_5(13) + q^6U_5(19)\\
&= \frac{(q;q)_\infty}{\left(q^{25};q^{25}\right)_\infty}\left(q^2\Sigma(5,-100,75) + q^{22}\Sigma(20,-25,75) - q^{66}\Sigma(50,50,75) - q^{116}\Sigma(50,125,75)\right)\\
&\quad + q\frac{(q;q)_\infty\left(q^{15};q^{15}\right)_\infty^3}{\left(q^5;q^5\right)_\infty\left(q^{25};q^{25}\right)_\infty}.
\end{aligned}
\tag{4.15}
$$

Similarly we reorder the $\Sigma(z,w,q)$ terms and apply Propositions 4.4 and 4.5 with $q \mapsto q^5$ to get that

$$\begin{aligned}
&q^2U_5(7) - q^3U_5(10) - q^4U_5(13) + q^5U_5(16)\\
&= q^{61}\Sigma(35,50,75) + q^{126}\Sigma(65,125,75) - q^3\Sigma(5,-100,75) - q^{23}\Sigma(20,-25,75)\\
&\quad + q^2\left(\Sigma(5,-115,75) + q^{25}\Sigma(20,-10,75) + q^{75}\Sigma(50,65,75) + q^{105}\Sigma(50,110,75)\right)\\
&\quad + q^5\left(\Sigma(5,-70,75) + q^{10}\Sigma(20,-55,75) + q^{35}\Sigma(35,5,75) + q^{160}\Sigma(65,170,75)\right)\\
&\quad - q^4\left(\Sigma(5,-85,75) + q^{15}\Sigma(20,-40,75) + q^{50}\Sigma(35,35,75) + q^{135}\Sigma(65,140,75)\right)\\
&\quad - q^{47}\left(\Sigma(35,20,75) + q^{40}\Sigma(50,80,75) + q^{50}\Sigma(50,95,75) + q^{105}\Sigma(65,155,75)\right)\\
&= \left(q^{60}\Sigma(35,50,75) + q^{125}\Sigma(65,125,75)\right)\left(-\frac{\left[q^{10};q^{15}\right]_\infty}{\left[q^5;q^{15}\right]_\infty} + q + q^2\frac{\left[q^5;q^{15}\right]_\infty}{\left[q^{10};q^{15}\right]_\infty}\right)\\
&\quad - \left(q^2\Sigma(5,-100,75) + q^{22}\Sigma(20,-25,75)\right)\left(-\frac{\left[q^{10};q^{15}\right]_\infty}{\left[q^5;q^{15}\right]_\infty} + q + q^2\frac{\left[q^5;q^{15}\right]_\infty}{\left[q^{10};q^{15}\right]_\infty}\right)\\
&\quad + q^2\frac{\left(q^{15};q^{15}\right)_\infty^3}{\left(q^5;q^5\right)_\infty} - q^2\frac{\left(q^{15};q^{15}\right)_\infty^3}{\left(q^5;q^5\right)_\infty}\\
&= \frac{(q;q)_\infty}{\left(q^{25};q^{25}\right)_\infty}\left(q^2\Sigma(5,-100,75) + q^{22}\Sigma(20,-25,75) - q^{60}\Sigma(35,50,75) - q^{125}\Sigma(65,125,75)\right).
\end{aligned} \tag{4.16}$$

Theorem 1.3 follows by equations (4.13), (4.15), and (4.16).

5. Proof of Theorem 1.4

Proof. Using Theorem 2.1 of [1] we find that

$$\begin{aligned}
\widetilde{\mathrm{PP}}(z,q) &= \sum_{n=1}^{\infty}\frac{q^n}{\left(zq^n;q\right)_\infty}\sum_{k=0}^{\infty}\frac{\left(zq;q\right)_k z^{-k}q^{nk}}{\left(q;q\right)_k}\\
&= \sum_{n=1}^{\infty}\sum_{k=0}^{\infty}\frac{z^{-k}q^{n+kn}}{\left(zq^{n+k};q\right)_\infty\left(q;q\right)_k}\\
&= \sum_{n=1}^{\infty}\frac{q^n}{\left(zq^n;q\right)_\infty} + \sum_{n=1}^{\infty}\sum_{k=1}^{\infty}\frac{z^{-k}q^{n+kn}}{\left(zq^{n+k};q\right)_\infty\left(q;q\right)_k}\\
&= \sum_{n=1}^{\infty}\frac{q^n}{\left(zq^n;q\right)_\infty} + \sum_{n=1}^{\infty}\sum_{k=1}^{\infty}\frac{q^n}{\left(q^n;q\right)_k\left(zq^{n+k};q\right)_\infty}\cdot\frac{z^{-k}q^{kn}\left(q;q\right)_{n+k-1}}{\left(q;q\right)_{n-1}\left(q;q\right)_k}.
\end{aligned} \tag{5.1}$$

The first term in (5.1) is the generating function for non-empty partitions with the power of q giving the number being partitioned and the power of z giving one less than the number of parts of the partition. This is the generating function for partition pairs from $\widetilde{\mathrm{PP}}$ where π_2 is empty, the power of q gives the number being partitioned and the power of z gives the paircrank.

In the second term in (5.1), we interpret the summands as follows. We have that $\frac{q^n}{\left(q^n;q\right)_k\left(zq^{n+k};q\right)_\infty}$ is the generating function for non-empty partitions π_1 where n is the smallest part, the power of q counts the number being partitioned, and the power of z counts the number of parts that are at least $n+k$. We have that $\frac{z^{-k}q^{kn}\left(q;q\right)_{n+k-1}}{\left(q;q\right)_{n-1}\left(q;q\right)_k}$ is the generating function for partitions π_2 with exactly k parts with each part between n and $2n-1$, where the power of q counts the number being partitioned

and the power of z counts the negative of the number of parts. Thus the second term in (5.1) is the generating function for partition pairs from $\widetilde{\mathrm{PP}}$, with both π_1 and π_2 non-empty, with the power of q giving the number being partitioned and the power of z giving the paircrank.

This proves that $C(m, n)$ is the number of partition pairs from $\widetilde{\mathrm{PP}}$ of n with paircrank m. These rearrangements and interpretations are essentially what was done in [8] as an intermediate step in getting an spt-crank defined on marked overpartitions. Also this is quite similar to the steps in [3] where Andrews and Garvan gave the ordinary partition crank after the vector partition crank. □

Acknowledgments

The author would like to thank Frank Garvan for suggesting this problem and for his help and encouragement.

References

[1] G. E. Andrews. *The theory of partitions*. Addison-Wesley Publishing Co., Reading, Mass.-London-Amsterdam, 1976. Encyclopedia of Mathematics and its Applications, Vol. 2.
[2] G. E. Andrews. The number of smallest parts in the partitions of n. *J. Reine Angew. Math.*, 624:133–142, 2008.
[3] G. E. Andrews and F. G. Garvan. Dyson's crank of a partition. *Bull. Amer. Math. Soc. (N.S.)*, 18(2):167–171, 1988.
[4] A. O. L. Atkin and P. Swinnerton-Dyer. Some properties of partitions. *Proc. London Math. Soc. (3)*, 4:84–106, 1954.
[5] K. Bringmann, J. Lovejoy, and R. Osburn. Rank and crank moments for overpartitions. *J. Number Theory*, 129(7):1758–1772, 2009.
[6] S. H. Chan. Generalized Lambert series identities. *Proc. London Math. Soc. (3)*, 91(3):598–622, 2005.
[7] A. B. Ekin. Some properties of partitions in terms of crank. *Trans. Amer. Math. Soc.*, 352(5):2145–2156, 2000.
[8] F. G. Garvan and C. Jennings-Shaffer. The spt-crank for overpartitions. *Acta Arith.*, 166(2):141–188, 2014.
[9] F. G. Garvan and H. Yesilyurt. Shifted and shiftless partition identities. II. *Int. J. Number Theory*, 3(1):43–84, 2007.
[10] J. Lovejoy and R. Osburn. Rank differences for overpartitions. *Q. J. Math.*, 59(2):257–273, 2008.
[11] J. Lovejoy and R. Osburn. M_2-rank differences for partitions without repeated odd parts. *J. Théor. Nombres Bordeaux*, 21(2):313–334, 2009.
[12] J. Lovejoy and R. Osburn. M_2-rank differences for overpartitions. *Acta Arith.*, 144(2):193–212, 2010.
[13] L. J. Slater. A new proof of Rogers's transformations of infinite series. *Proc. London Math. Soc. (2)*, 53:460–475, 1951.

Department of Mathematics, University of Florida, Gainesville, Florida 32611, USA
cjenningsshaffer@ufl.edu